



MAMMALIAN TOXICOLOGY OF DIMETHYL AMIDES

Applicable to these current Stepan products:

STEPOSOL® M-10 HALLCOMID® M-8-10	STEPOSOL® M-8-10 NINOL® CAA	HALLCOMID® M-10
-------------------------------------	--------------------------------	-----------------

Toxicological Information:

<u>Test/Conditions</u>	<u>Results/Classification</u>	<u>References</u>
Acute Oral Toxicity (rat)(gavage)(14 days) n=5/sex/dose	LD ₅₀ = 1750 mg/kg (slightly toxic orally)	Stepan Study No. 90-023A
Acute Dermal Toxicity (rat)(dermal application)(15 days) n=5/sex/dose	LD ₅₀ = 2000 mg/kg (slightly toxic dermally)	Stepan Study No. 95-029A
Bovine Corneal Opacity Permeability Test (BCOP)	18.7 - 26.6 (mild to moderate eye irritant)	Stepan Study No. 08-005A; 11-003A
Sensitization Study (guinea pig)(6 hr. exposure) n=46	Not a skin sensitizer	Stepan Study No. 90-020A
Genotoxicity Study (Ames)(bacteria)(48 hr)	Non-mutagenic	Stepan Study No. 92-030A
Genotoxicity Study (chromosomal aberration)(4 hr exposure)	Not considered to be clastogenic	Stepan Study No. 95-028A
Genotoxicity Study (unscheduled DNA synthesis)(24 hrs)	Negative	Stepan Study No. 91-049A
Reproductive Toxicity (rat)(90 day study)	NOAEL= 10,000 ppm	US HPV

References:

STEPOSOL® ; HALLCOMID® ; NINOL® are registered trademarks of Stepan Company.

Last Update: 6.28.11

Revision Reference: TX056-02

Nothing contained herein grants or extends a license, express or implied, in connection with patents, issued or pending, of the manufacturer or others. The information contained herein is based on the manufacturer's own study and the works of others. The manufacturer makes no warranties, expressed or implied, as to the accuracy, completeness, or adequacy of the information contained herein. The manufacturer shall not be liable (regardless of fault) to the vendee's employees, or anyone for any direct, special or consequential damages arising out of or in connection with the accuracy, completeness, adequacy or furnishing of such information.